

# New data on sponges from Svalbard Archipelago with a description of a new species of *Halicnemida*

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

---

## Abstract

© 2018 Informa UK Limited, trading as Taylor & Francis Group. Sponge fauna from the west and south of Svalbard archipelago was studied. A total of 28 species of sponges were identified. Five of them are new records to the study area and one is new to science. New records for Spitsbergen are: *Axinella hispida* (Koltun 1959), *Haliclona rossica* (Hentschel 1929), *Myxilla* (*Myxilla*) *perspinosa* (Lundbeck 1905), *Sphaerotylus borealis* (Swartschewsky 1906) and *Suberites carnosus* (Johnston 1842). A new species of sponge, *Halicnemida wagini* sp. nov., collected south-west of Spitsbergen on the continental slope at a depth of 423–425 m, is described. It is similar to *Halicnemida patera* (Bowerbank 1864) and *H. verticillata* (Bowerbank 1866) in skeleton architecture, but significantly differs in its spicule complement and external appearance. Primarily, distinct differences concern the morphology of microscleres. In comparison with *Halicnemida patera*, the new species has an additional category of acanthose microscleres presented by microstrongyles, while microscleres of *H. verticillata* are only slightly curved and also verticillately spined. The study area, west of Spitsbergen, establishes close relationships with North Atlantic due to warm Atlantic waters of the West Spitsbergen Current. Proximity of the new species to the North Atlantic representatives of *Halicnemida* genus allowed us to suggest that *H. wagini* sp. nov., both ecologically and morphologically, has stood apart as a distinct biological species in the conditions of the Arctic. <http://www.zoobank.org/lsid:zoobank.org:pub:2A09537B-63D7-4F07-A2AB-9C707A686333>.

<http://dx.doi.org/10.1080/00222933.2018.1440020>

---

## Keywords

Arctic, *Halicnemida*, new species, Porifera, Stelligeridae, Svalbard Archipelago

## References

- [1] Anisimova NA, Jørgensen LL, Lyubin PA, Manushin IE., 2011. The Barents Sea-ecosystem, resources, management: half a century of Russian-Norwegian cooperation. Trondheim: Tapir Academic Press. Chapter Benthos; p. 121–159.
- [2] Blindheim J, Østerhus S. 2005. The Nordic Seas, main oceanographic features. In: Drange H, Dokken T, Furevik T, Gerdes R, Berger W, editors. The Nordic Seas: an integrated perspective. Washington (DC): American Geophysical Union; p. 11–37.
- [3] Bluhm BA, Ambrose W, Bergmann M, Clough L, Gebruk A, Hasemann C, Iken K, Klages M, MacDonald I, Renaud P, et al. 2011. Diversity of the arctic deep-sea benthos. *Marine Biodiversity*. 41:87–107.
- [4] Bowerbank JS. 1864. A monograph of the British spongiadae. Volume 1. *Ray Soc.* 36:1–290.
- [5] Bowerbank JS. 1866. A monograph of the British spongiadae. Volume 2. *Ray Soc.* 38:1–388.
- [6] Bowerbank JS. 1874. A monograph of the British spongiadae. Volume 3. *Ray Soc.* 51:1–367.

- [7] Breitung L. 1896. Kalkschwämme der Bremer Expedition nach Ost-Spitzbergen im Jahre 1889 (Prof. W. Küenthal und Dr. A. Walter). *Zool Anz.* 19(513):426–432.
- [8] Buzhinskaja GN, editor. 2011. Illustrated keys to free-living invertebrates of Eurasian arctic seas and adjacent deep waters: volume 2. Fairbanks: University of Alaska Fairbanks.
- [9] Cárdenas P, Pérez T, Boury-Esnault N. 2012. Sponge systematics facing new challenges. *Adv Sponge Sci Phyl Syst Ecol.* 61:79–209.
- [10] Cárdenas P, Rapp HT. 2012. A review of Norwegian streptaster-bearing Astrophorida (Porifera: Demospongiae: Tetractinellida), new records and a new species. *Zootaxa.* 3253:1–52.
- [11] Carter HJ. 1875. Notes introductory to the study and classification of the Spongida. Part II. Proposed classification of the Spongida. *Ann Magazine Nat Hist.* 4(16):126–145.
- [12] Dendy A. 1922. Reports of the Percy Sladen Trust expedition to the Indian Ocean in 1905, Volume 7: report on the Sigmatotetraxonida collected by H.M.S. 'Sealark' in the Indian Ocean. *Trans Linnean Soc London.* 18(1):1–164, pls 1–18.
- [13] Dunton K. 1992. Arctic biogeography: the paradox of the marine benthic fauna and flora. *Trends Ecol Evol.* 7:183–189.
- [14] Fristedt K. 1887. Sponges from the Atlantic and Arctic Oceans and the Behring Sea. *Vega-Expeditionens Vetenskap lakttagelser (Nordenskiöld).* 4:401–471.
- [15] Golikov AN, Scarlato OA. 1989. Evolution of Arctic ecosystems during the Neogene period. In: Herman Y, editor. *The Arctic seas: climatology, oceanography, geology, and biology.* New York (NY): Springer; p. 257–279.
- [16] Grant RE. 1841. *Outlines of comparative anatomy.* London: Bailliere; p. 1–656.
- [17] Gray JE. 1867. Notes on the arrangement of Sponges, with the descriptions of some New Genera. *Proc Zool Soc London.* 2:492–558. pls XXVII–XXVIII.
- [18] Hentschel E. 1916. 3- Die Spongien des Eisfjords (In: *Zoologische Ergebnisse der Schwedischen Expedition nach Spitzbergen, 1908. Teil II.*). *K Sven Vetenskaps Handlingar.* 54(3):1–18.
- [19] Hentschel E. 1929. Die Kiesel- und Hornschwämme des Nördlichen Eismeers. In: Römer F, Schaudinn F, Brauer A, Arndt W, editors. *Fauna Arctica. Eine Zusammenstellung der arktischen Tierformen mit besonderer Berücksichtigung des Spitzbergen-Gebietes auf Grund der Ergebnisse der Deutschen Expedition in das Nördliche Eismeer im Jahre 1898.* Vol. 5(4). Jena: G. Fischer; p. 857–1042. pls XII–XIV.
- [20] Higgin T. 1877. Description of some Sponges obtained during a Cruise of the Steam-Yacht 'Argo' in the Caribbean and neighbouring Seas. *Ann Magazine Nat Hist.* 19(4):291–299. pl. XIV.
- [21] Johnston G. 1842. *A history of British sponges and lithophytes.* Edinburgh: WH Lizars; p. 1–264, pls. I–XXV.
- [22] Koltun VM. 1959. Кремнегоровые губки северных и дальневосточных морей СССР [Siliceous horny sponges of the northern and far-eastern seas of the U.S.S.R.]. In: Pavlovsky EN, editor. *Identifiers of the USSR fauna issued by the Zoological Institute of the Academy of Sciences of the USSR.* Moscow–Leningrad: Nauka, Academy of Sciences of the USSR; Vol. 67, p. 1–263. Russian.
- [23] Koltun VM. 1966. Четырехлучевые губки северных и дальневосточных морей СССР [Four rayed sponges (order Tetraxonida) of the Northern and Far-Eastern Seas of the USSR]. In: Bychowski BE, editor. *Identifiers of the USSR fauna issued by the Zoological Institute of the Academy of Sciences of the USSR.* Moscow–Leningrad: Nauka, Academy of Sciences of the USSR; Vol. 90, p. 1–112. Russian.
- [24] Koltun VM. 1967. Стекланные или Шестилучевые, губки северных и дальневосточных морей СССР [Glass, or Hexactinellid sponges of the Northern and Far-Eastern Seas of the USSR (Class Hyalospongiae)]. In: Bychowski BE, editor. *Identifiers of the USSR fauna issued by the Zoological Institute of the Academy of Sciences of the USSR.* Moscow–Leningrad: Nauka, Academy of Sciences of the USSR; Vol. 94, p. 1–124. Russian.
- [25] Lévi C. 1953. Sur une nouvelle classification des Démosponges. *C R Hebd Séances l'Acad Sci Paris.* 236(8):853–855.
- [26] Lundbeck W. 1905. Porifera (Part II.) Desmacidonidae (pars.). In: Hagerup H, editor. *The Danish ingolf-expedition.* Vol. 6. Copenhagen: BiancoLuno; Vol. 6(2), p. 1–219.
- [27] Mironov AN, Dilman AB. 2008. Effect of the East Siberian barrier on the echinoderm dispersal in the Arctic Ocean. *Oceanology.* 50(3):342–355.
- [28] Morrow C, Cárdenas P. 2015. Proposal for a revised classification of the Demospongiae (Porifera). *Front Zool.* 12:7.
- [29] Morrow C, Picton B, Erpenbeck D, Boury-Esnault N, Maggs C, Allcock A. 2012. Congruence between nuclear and mitochondrial genes in Demospongiae: A new hypothesis for relationships within the G4 clade (Porifera: Demospongiae). *Mol Phylogenet Evol.* 62(1):174–190.
- [30] Morrow CC, Redmond NE, Picton BE, Thacker RW, Collins AG, Maggs CA, Sigwar JD, Allcock AL. 2013. Molecular phylogenies support homoplasy of multiple morphological characters used in the taxonomy of Heteroscleromorpha (Porifera: Demospongiae). *Integr Comp Biol.* 53:428–446.
- [31] Nardo GD. 1833. Auszug aus einem neuen System der Spongiarien, wonach bereits die Aufstellung in der Universitäts-Sammlung zu Padua gemacht ist. *Isis Oken.* 519–523.

- [32] Nesis KN. 1983. A hypothesis of the origin of Western and Eastern Arctic ranges of marine bottom animals. *Russ J Marine Biol.* 5:3-13.
- [33] Plotkin A. 2004. Biodiversity and distribution of Polymastiidae (Demospongiae, Hadromerida) in the Arctic area. *Boll Mus Ist Biol Univ Genova.* 68:535-547.
- [34] Plotkin A, Gerasimova E, Rapp HT. 2017. Polymastiidae (Porifera: Demospongiae) of the Nordic and Siberian Seas. *J Marine Biol Assoc U K.* 1-63.
- [35] Plotkin A, Voigt O, Willassen E, Rapp HT. 2016. Molecular phylogenies challenge the classification of Polymastiidae (Porifera, Demospongiae) based on morphology. *Org Divers Evol.* 17(1):45-66.
- [36] Saloranta TM, Svendsen H. 2001. Across the Arctic Front west of Spitsbergen: high-resolution CTD sections from 1998-2000. *Polar Res.* 20:177-184.
- [37] Sarà M, Bavestrello G, Mensi P. 1992. Redescription of *Tethya norvegica* Bowerbank (Porifera, Demospongiae), with remarks on the genus *Tethya* in the North East Atlantic. *Zool Scr.* 21(3):211-216.
- [38] Schmidt O. 1862. Die Spongien des adriatischen Meeres. Leipzig: WilhelmEngelmann; p. i-viii, 1-88, pls 1-7.
- [39] Schmidt O. 1870. Grundzüge einer Spongien-Fauna des atlantischen Gebietes. Leipzig: WilhelmEngelmann; p. iii-iv, 1-88, pls I-VI.
- [40] Sim CJ, Kim HY, Byeon HS. 1990. A systematic study on the marine sponges in Korea 8. Tetractinomorpha. *Korean J Syst Zool.* 6(1):123-144.
- [41] Sollas WJ. 1885. A classification of the sponges. *Ann Magazine Nat Hist.* 16(95):395.
- [42] Steenstrup E, Tendal OS. 1982. The genus *Thenea* (Porifera, Demospongia, Choristida) in the Norwegian sea and adjacent waters; an annotated key. *Sarsia.* 67(4):259-268.
- [43] Swartschewsky BA. 1906. Beiträge zur Spongien-Fauna des Weissen Meeres. *Mém Soci Nat Kiev.* 20:307-371.
- [44] Teigen SH., 2011. Water mass exchange in the sea west of Svalbard [dissertation]. Bergen, Norway: University of Bergen.
- [45] Topsent E. 1897. Sur le genre *Halicnemida* Bowerbank. *Mém Soc Zool France.* 10:235-251.
- [46] Topsent E. 1898. Eponges nouvelles des Açores. (Première Serie) *Mém Soc Zool France.* 11:225-255.
- [47] Topsent E. 1928. Spongiaires de l'Atlantique et de la Méditerranée provenant des croisières du Prince Albert Ier de Monaco. Résultats des campagnes scientifiques accomplies par le Prince Albert I. Monaco. 74:1-376, pls I-XI.
- [48] Van Soest RWM. 1987. Biogeographic and taxonomic notes on some Eastern Atlantic sponges. In: Jones WC, editor. *European Contributions to the Taxonomy of Sponges.* Sherkin Island, County Cork: Sherkin Island Marine Station; p. 13-28.
- [49] Van Soest RWM, Boury-Esnault N, Hooper JNA, Rützler K, De Voogd NJ, Alvarez B, Hajdu E, Pisera AB, Manconi R, Schönberg C, et al. 2017. World Porifera database [accessed 2017 Dec]. <http://www.marinespecies.org/porifera>
- [50] Van Soest RWM. 2017. *Halicnemida* Bowerbank, 1864 [cited 2016 Dec]. In: Van Soest RWM, Boury-Esnault N, Hooper JNA, Rützler K, De Voogd NJ, Alvarez De Glasby B, Hajdu E, Pisera AB, Manconi R, Schoenberg C, et al., editors. *World Porifera database.* <http://www.marinespecies.org/porifera/porifera.php?p=taxdetails&id=131790>.
- [51] Von Lendenfeld R. 1898. Die Clavulina der Adria. *Nova Acta Acad Caes Leop Carol German Nat Cur.* 69:1-251, pls I-XII.
- [52] Walczowski W. 2013. Frontal structures in the West Spitsbergen Current margins. *Ocean Sci.* 9:957-975.
- [53] Walczowski W, Piechura J. 2007. Pathways of the Greenland Sea warming. *Geophys Res Lett.* 34(10):L10608.